

# **NOTICE:**

**“BEST AVAILABLE COPY”**

**PORTIONS OF THE FOLLOWING  
DOCUMENT ARE ILLEGIBLE**

**The Administrative Record Staff**



## ESTIMATING SOIL MIXING BY RODENTS

E M White and D C Carlson  
Plant Science Dept (Soils)  
South Dakota State University  
Brookings, South Dakota 57007

## ABSTRACT

The importance of soil mixing by prairie dogs (*Cynomys* sp) can be estimated by observation and quantification. Rodent burrows, which have been filled with dark-colored surface soil, can be observed in subsoil exposed by construction activities or by natural erosion along gullies. Burrow entrances surrounded by soil mounds are easily located, particularly for prairie dogs. Prairie dogs have from 25 to 125 mounds per ha with 62 as an average number. With this average number, a one ha area would contain 18 m<sup>2</sup> of mounds if their diameter is 0.6 m. If new mounds were constructed at new locations each year, about 550 years would be needed to cover the entire ha with mounds. Sixty-two 15-cm diameter burrows would have an area of 1.1 m<sup>2</sup> so that 8,800 years would be needed for a ha of burrows to be created. Obviously, new burrows are not constructed each year. These calculations indicate that the effect of soil mixing by rodents may be exaggerated in the literature.

## INTRODUCTION

Soil mixing by rodents is important in soil genesis because unweathered parent materials are moved to the surface where weathering is more rapid. Several species of rodents have been reported to have a significant effect on soil development because of their burrowing activities (Joffe 1949). Nearly a complete mixing of soil has been reported in prairie dog (*Cynomys* sp) towns (Thorp 1949). Koford (1958) discussed the general relationships of prairie dogs to soils. Burrow systems have been traced and described for black-tailed prairie dogs (*C. ludovicianus*) in South Dakota (Sheets et al 1971).

Green (1960) reported *Cynomys* fossils in Tertiary strata in South Dakota so the species may have been present intermittently since that time. Because prairie dog bones do occur in archaeological soils, prairie dogs likely have been present during the Holocene epoch and probably in interglacial intervals during the Pleistocene. If prairie dogs have occurred in these soils during most of the Pleistocene, more evidence of soil mixing should have been reported in the literature. The lack of evidence may be due to the fact that most studies are based on observations of surface mounds of soil dug from burrows without considering the probability that a soil profile has been altered. This paper will consider this probability for prairie dogs.

The number of burrows/ha (25/acre) proportional to the frequency of burrows/ha (25/acre) (Koford 1958), but F. Lechleitner (1966) found 224, 419, and 34 burrows/acre, a scatter of burrows or mound gully or trench area would likely occur in a 200 x 200 m area (200 x 200 m area). A model intercept model estimate the frequency of burrows, respectively. A second transect first to form the side of the transect line rows were used, but quadratic equation, crossed the circle formed were calculated of the circles inside the geometric relationship lines.

The frequency of burrows found to occur along a transect line and 1070 m, respectively. The average transect line distances were 24.6 cm and 0.8 cm, respectively. Soil in 62 61-cm diameter burrows would be needed to cover a 1 ha area. Sixty-two 15-cm-diameter burrows would be needed to cover a 1 ha area. Eight thousand burrows across an area of 1 ha. The average is to divide the transect line by the frequency of burrows respectively. If an average mound

RECORD

1984

## RODENTS

on  
)  
ity  
007

prairie dogs (*Cynomys* sp.)  
Rodent burrowed surface soil, can  
activities or by  
surroundings  
prairie dogs  
with 62 as an average  
area would contain  
If new mounds were  
550 years would be  
Sixty-two 15-cm  
m<sup>2</sup> so that 8,800 years  
Obviously, new  
calculations indi-  
cations may be exaggerated

in soil genesis because  
to the surface where  
of rodents have been  
development because  
Nearly a complete mix-  
og (*Cynomys* sp.) towns  
he general relationships  
have been traced and  
(*Ludovicianus*) in South

ils in Tertiary strata in  
en present intermittently  
do occur in archaeological  
sents during the Holocene  
ils during the Pleistocene  
soils during most of the  
should have been reported  
ay be due to the fact that  
of surface mounds of  
he probability that a  
l consider this probability

## METHODS

The number of prairie dog burrows in an area is generally proportional to the forage available for grazing, with about 62 burrows/ha (25/acre) being an average number according to Koford (1958), but Fitzgerald and Lechleitner (1974), Tileston and Lechleitner (1966), and Bishop and Culbertson (1976), reported 224, 419, and 34 mounds per acre. Based on an average of 25 burrows/acre, a scaled model was used to estimate the number of burrows or mounds that would be exposed along the edge of a gully or trench across an area and the amount of soil mixing that would likely occur. The model had 25 burrows arrayed in a 61 x 61 m area (200 x 200 ft with burrows located at ordinate and abscissa values of 20, 60, 100, 140, and 180 ft). Line transects across the model intercept mounds and burrows with a frequency that should estimate the frequency with which a gully or trench would intercept them. Sixty and 150 transects were used for mounds and burrows, respectively, in the model. Ends of transects across the model were located from random number tables (0 to 200 range). A second transect was located 76 cm (30 inches) away from the first to form the sides of a belt transect across the area. Equations of the transect lines and the circles representing mounds or burrows were used, by substitution and solution of the resulting quadratic equation, to calculate the points where transect lines crossed the circle circumferences. The lengths of the chords formed were calculated from these line-circle intercepts. Areas of the circles inside the belt transects also were determined from the geometric relationships of the circles and the two parallel transect lines.

## RESULTS

The frequency with which a single mound and burrow was found to occur along a line transect was one in a length of 191 and 1070 m, respectively. With the 76 cm wide belt transect, the respective distances were 85 m and 209 m. The mean lengths of the average transect that were in a circular mound or burrow, i.e., 24.6 cm and 0.8 cm, are very small.

Soil in 62 61-cm-diameter mounds would have an area of 11 m<sup>2</sup>/ha. If a new mound were constructed each year, 550 yrs would be needed to cover the entire surface hectare with mounds. Sixty-two 15-cm-diameter burrows would have an area of 1.1 m<sup>2</sup>/ha. Eight thousand eight hundred yrs would be needed to form burrows across an area. Another procedure for estimating surface area is to divide mean chord lengths for mounds and burrows by the transect lengths. These values are 261 and 7,860 years, respectively.

If an average mound has a diameter of 0.6 m and is 0.3 m high,

2

the volume of soil present would be 0.023 m<sup>3</sup> after the 15-cm-diameter burrow is subtracted. A hectare of one meter thick soil has a volume of 10,000 m<sup>3</sup> and 62 mounds/ha would have a volume of 1.4 m<sup>3</sup>/ha. More than 7,000 yrs would be needed to invert the subsoil and surface soil. This estimate neglects the difference in the bulk densities of the surface and subsoil, the construction of mounds within older mounds, and the length of time a burrow is used before a new one is constructed.

#### DISCUSSION

Soil profiles exposed along the banks of gullies and roads contain an occasional burrow, but the overall movement of subsoil to surface layers appears to be minor. Dark-colored surface soil that infiltrates into burrows is very evident in the light-colored subsoil. Although the organic matter, which causes the darkening, may be destroyed by oxidation, archaeologists find dark-colored infills in holes dug by prehistoric inhabitants that are several thousand years old. Floodplain sediments, which accumulated at a slow rate, contain horizontally bedded layers which differ in texture and in the content of dark organic matter. These sediments have little evidence of soil mixing by rodents. From observations (White, 1975), Tertiary-age soft siltstone frequently contains some thin more consolidated but fractured laminae that remain in place after soils have formed through them. It would not be possible to identify these laminae if rodents had mixed the soil. Stratified fluvial gravels exposed in banks of gravel pits also contain little evidence of mixing of the strata although some of the gravels were deposited hundreds of thousands of years ago.

No mechanism apparently has been proposed that would cause heterogeneous materials to be reorganized into strata in subsoils. Our calculations indicate that a long time is needed for the surface soil and subsoil to be completely mixed. Based on liberal assumptions, more than 7,000 yrs would be needed for prairie dogs to move all subsoil to the surface of an area. This is a minimum estimate which does not allow mixing of soil twice in the period. Thus, the logical conclusions are that soil mixing by prairie dogs is not a major process in soil genesis and that rodents have not caused extensive soil mixing in South Dakota.

#### LITERATURE CITED

- Bishop, N. G. and J. L. Culbertson. 1976. Decline of prairie dog towns in southwestern North Dakota. *J. Range Manage.* 29:217-220.
- Fitzgerald, J. P. and R. R. Lechleitner. 1974. Observations on the biology of Gunnison's prairie dog in central Colorado. *Amer. Midl. Nat.* 92:163.
- Green, M. 1960. A Tertiary *Cynomys* from South Dakota. *J. Paleont.* 34:547.

Joffe, J  
Koford, Mon  
Sheets, F prair  
Thorp, J 68 18  
Tilston, tailec Midl  
White, E Consc

- Joffe, J S 1949 *Pedology* Pedology Publications New Brunswick, N Y
- Koford, C B 1958 Prairie dogs, white faces, and blue grama *Wildlife Monogr* 3 1-78
- Sheets, R G, R L Linder, and R B Dahlgren 1971 Burrow systems of prairie dogs in South Dakota *J Mammal* 52 451-453
- Thorp, J 1949 Effects of certain animals that live in soils *Sci Monthly* 68 180-191
- Tileston, J V and R R Lechleitner 1966 Some comparisons of the black-tailed and white-tailed prairie dog in north central Colorado *Amer Midl Nat* 75 292-316
- White, E M 1975 Soil Survey of Mellette County, South Dakota Soil Conservation Service, U S D A

5-cm-diameter soil has a volume of vert the sub-  
erence in the  
struction of  
a burrow is

id roads con-  
of subsoil to  
face soil that  
colored subsoil  
rkening, may  
colored infills  
ral thousand  
d at a slow  
er in texture  
diments have  
utions (White,  
is some thin  
ain in place  
ot be possible  
oil Stratified  
contain little  
f the gravels

t would cause  
a in subsoils  
l for the sur-  
on liberal as-  
r prairie dogs  
s a minimum  
in the period  
y prairie dogs  
nts have not

ie dog towns in  
20

on the biology  
idl Nat 92 146-

Paleont 34 545-

14